

Refine Search

Search Results -

Terms	Documents
L17 and L16	3

Database:

US Pre-Grant Publication Full-Text Database
US Patents Full-Text Database
US OCR Full-Text Database
EPO Abstracts Database
JPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Search:

L18

Refine Search

Recall Text

Clear

Interrupt

Search History

DATE: Wednesday, December 21, 2005 [Printable Copy](#) [Create Case](#)

Set Name Query

side by side

Hit Count

Set Name

result set

DB=PGPB,USPT,USOC,EPAB,JPAB,DWPI,TDBD; PLUR=YES; OP=OR

<u>L18</u>	L17 and l16	3	<u>L18</u>
<u>L17</u>	halbert.in.	797	<u>L17</u>
<u>L16</u>	L15 and estradiol	2618	<u>L16</u>
<u>L15</u>	L14 and PCMA cholesteryl Oleate	70787	<u>L15</u>
<u>L14</u>	L13 and perfluorinated cholesteryl oleate	70787	<u>L14</u>
<u>L13</u>	L12 and bishydroxycoumarin	44	<u>L13</u>
<u>L12</u>	L11 and methotrexate diester	84375	<u>L12</u>
<u>L11</u>	L10 and retinoic acid	2531878	<u>L11</u>
<u>L10</u>	L9 and cholesteryl oleate	69883	<u>L10</u>
<u>L9</u>	L7 and fatty acid	2531879	<u>L9</u>
<u>L8</u>	L7 and retinyl derived compounds	3721543	<u>L8</u>
<u>L7</u>	L6 and pyrenes	5739	<u>L7</u>
<u>L6</u>	L5 and (polyunsaturated compounds)	512666	<u>L6</u>
<u>L5</u>	L4 and (Apo B binding site)	642865	<u>L5</u>
<u>L4</u>	L3 and cholesterol ester	901490	<u>L4</u>
<u>L3</u>	L2 and lipophilic substituent	356677	<u>L3</u>
<u>L2</u>	LDL and liposome	3716	<u>L2</u>
<u>L1</u>	LDL and liposome	3716	<u>L1</u>

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1653HXP

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 SEP 09 ACD predicted properties enhanced in REGISTRY/ZREGISTRY
NEWS 4 OCT 03 MATHDI removed from STN
NEWS 5 OCT 04 CA/CAPLUS-Canadian Intellectual Property Office (CIPO) added
to core patent offices
NEWS 6 OCT 13 New CAS Information Use Policies Effective October 17, 2005
NEWS 7 OCT 17 STN(R) AnaVist(TM), Version 1.01, allows the export/download
of CAPLUS documents for use in third-party analysis and
visualization tools
NEWS 8 OCT 27 Free KWIC format extended in full-text databases
NEWS 9 OCT 27 DIOGENES content streamlined
NEWS 10 OCT 27 EPFULL enhanced with additional content
NEWS 11 NOV 14 CA/CAPLUS - Expanded coverage of German academic research
NEWS 12 NOV 30 REGISTRY/ZREGISTRY on STN(R) enhanced with experimental
spectral property data
NEWS 13 DEC 05 CASREACT(R) - Over 10 million reactions available
NEWS 14 DEC 14 2006 MeSH terms loaded in MEDLINE/LMEDLINE
NEWS 15 DEC 14 2006 MeSH terms loaded for MEDLINE file segment of TOXCENTER
NEWS 16 DEC 14 CA/CAPLUS to be enhanced with updated IPC codes
NEWS 17 DEC 16 MARPATprev will be removed from STN on December 31, 2005
NEWS 18 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the
IPC reform

NEWS EXPRESS DECEMBER 02 CURRENT VERSION FOR WINDOWS IS V8.01,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 02 DECEMBER 2005.
V8.0 USERS CAN OBTAIN THE UPGRADE TO V8.01 AT
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:25:53 ON 21 DEC 2005

=> file medline, biosis, fsta, jicst, wpids, biotechds		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.84	0.84

FILE 'MEDLINE' ENTERED AT 15:28:01 ON 21 DEC 2005

FILE 'BIOSIS' ENTERED AT 15:28:01 ON 21 DEC 2005

Copyright (c) 2005 The Thomson Corporation

FILE 'FSTA' ENTERED AT 15:28:01 ON 21 DEC 2005

COPYRIGHT (C) 2005 International Food Information Service

FILE 'JICST-EPLUS' ENTERED AT 15:28:01 ON 21 DEC 2005

COPYRIGHT (C) 2005 Japan Science and Technology Agency (JST)

FILE 'WPIDS' ENTERED AT 15:28:01 ON 21 DEC 2005

COPYRIGHT (C) 2005 THE THOMSON CORPORATION

FILE 'BIOTECHDS' ENTERED AT 15:28:01 ON 21 DEC 2005

COPYRIGHT (C) 2005 THE THOMSON CORPORATION

=> s LDL and liposome
L1 254 LDL AND LIPOSOME

=> s l1 and Apo b
L2 11 L1 AND APO B

=> d l2 ti abs ibib tot

L2 ANSWER 1 OF 11 MEDLINE on STN

TI Evaluation of antioxidant and prooxidant activities of bamboo

Phyllostachys nigra var. Henonis leaf extract in vitro.

AB Solvent-extracted bamboo leaf extract (BLE) containing chlorogenic acid, caffeic acid, and luteolin 7-glucoside was evaluated in vitro for free radical scavenging and antioxidant activities using a battery of test methods. BLE exhibited a concentration-dependent scavenging activity of DPPH radical. BLE prolonged the lag phase and suppressed the rate of propagation of **liposome** peroxidation initiated by peroxy radical induced by 2,2'-azobis(2-amidinopropane dihydrochloride (AAPH) at 37 degrees C. BLE also prevented human low-density lipoprotein oxidation, mediated by Cu(2+), which was monitored by the lower formation of conjugated diene and fluorescence and a reduced negative charge of **apo-B** protein. Finally, BLE protected supercoiled DNA strand against scission induced by AAPH-mediated peroxy radical. Prooxidant activity of BLE was seen in a Cu(2+)-induced peroxidation of structured phosphatidylcholine **liposome**, indicating catalytic peroxidation due to a relatively high reducing power of BLE. It was concluded that the BLE has both antioxidant activity and prooxidant activity; the antioxidant activity was attributed to free radical scavenging activity, and the prooxidant activity, albeit minor, resulted from the reducing power of plant phenolics in the presence of transitional metal ions.

ACCESSION NUMBER: 2000458826 MEDLINE

DOCUMENT NUMBER: PubMed ID: 10956087

TITLE: Evaluation of antioxidant and prooxidant activities of bamboo Phyllostachys nigra var. Henonis leaf extract in vitro.

AUTHOR: Hu C; Zhang Y; Kitts D D

CORPORATE SOURCE: Food, Nutrition and Health, Faculty of Agricultural Science, University of British Columbia, Vancouver, BC,

Canada.
SOURCE: Journal of agricultural and food chemistry, (2000 Aug) 48
(8) 3170-6.
Journal code: 0374755. ISSN: 0021-8561.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200009
ENTRY DATE: Entered STN: 20001005
Last Updated on STN: 20001005
Entered Medline: 20000925

L2 ANSWER 2 OF 11 MEDLINE on STN

TI Oxidative interaction of unpaired hemoglobin chains with lipids and
proteins: a key for modified serum lipoproteins in thalassemia.
AB We searched for a biochemical explanation to the modification of
lipoproteins like low-density lipoproteins (**LDL**) observed in
patients with the severe hemolytic anemia beta-thalassemia. Because a
large fraction of the **LDL** surface is composed of phospholipids,
we first explored the possible involvement of phospholipids in the
oxidative interaction of **LDL** with hemoglobin (Hb), using brain
extract phospholipid liposomes as a model. The relative binding affinity
and oxidative interaction of three hemoglobin variants (intact Hb A and
isolated beta- and alpha-chains) with **LDL** and **liposome**
were compared. Studies carried out at low pH/ionic strength and under
physiological conditions revealed that association of hemoglobin variants
with the phospholipid liposomes is driven by electrostatic forces but
their binding is not a prerequisite for oxidative interaction. Unlike
phospholipid liposomes, **LDL** underwent only a negligible
association with the Hb variants under all pH/ionic strength conditions.
Nevertheless, **LDL** induced oxidation of Hb variants, mostly
alpha-chains. The dissimilar behavior of the liposomes and **LDL**
indicated that **LDL** protein **apo B** rather than
phospholipids is the actual **LDL** surface component which
interacts with the hemoglobin variants. This agrees with the finding that
apo B protein underwent oxidative crosslinking by the
hemoglobin variants among which alpha-chains were most active. We
concluded from these results that the ability of hemoglobin to undergo
autooxidation is the key to its oxidative reactivity toward **LDL**.
The results of the present study indicate that the modified **LDL**
particles observed in beta-thalassemia may reflect lipoprotein oxidation
by alpha-chains in circulation.

ACCESSION NUMBER: 97428182 MEDLINE
DOCUMENT NUMBER: PubMed ID: 9281309
TITLE: Oxidative interaction of unpaired hemoglobin chains with
lipids and proteins: a key for modified serum lipoproteins
in thalassemia.
AUTHOR: Altamentova S M; Marva E; Shaklai N
CORPORATE SOURCE: Sackler Faculty of Medicine, Tel-Aviv University, Tel-Aviv,
698887, Israel.
SOURCE: Archives of biochemistry and biophysics, (1997 Sep 1) 345
(1) 39-46.
Journal code: 0372430. ISSN: 0003-9861.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199710
ENTRY DATE: Entered STN: 19971013
Last Updated on STN: 19971013
Entered Medline: 19971002

L2 ANSWER 3 OF 11 MEDLINE on STN
TI Low-density lipoproteins interact with **liposome**-binding sites on the cell surface.
AB Under physiological conditions significant amounts of low-density lipoprotein **LDL** particles are taken up by cells independently of specific high-affinity **LDL** receptors (**apo-B** receptors). Previously it was established that some cells contain surface sites capable of binding liposomes. We proposed that **liposome**-binding sites could contribute to **LDL** interaction with the cell surface via phospholipid molecules of **LDL** particles. To check this hypothesis we studied the competitive interaction of human **LDL** and DPPC liposomes with mouse embryo fibroblasts depleted of **apo-B** receptors by preliminary incubation with **LDL**. We have found that after removal of the **liposome**-binding sites from cell lamellae these areas of the cell surface lose their ability to bind **LDL**.

ACCESSION NUMBER: 91348212 MEDLINE
DOCUMENT NUMBER: PubMed ID: 1879530
TITLE: Low-density lipoproteins interact with **liposome**-binding sites on the cell surface.
AUTHOR: Galkina S I; Ivanov V V; Preobrazhensky S N; Margolis L B; Bergelson L D
CORPORATE SOURCE: Belozersky Laboratory of Molecular Biology and Bioorganic Chemistry, Moscow State University, USSR.
SOURCE: FEBS letters, (1991 Aug 5) 287 (1-2) 19-22.
Journal code: 0155157. ISSN: 0014-5793.
PUB. COUNTRY: Netherlands
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199109
ENTRY DATE: Entered STN: 19911020
Last Updated on STN: 19911020
Entered Medline: 19910927

L2 ANSWER 4 OF 11 MEDLINE on STN
TI [Apolipoprotein B of plasma lipoproteins incorporated in liposomes: immunological properties and organ distribution when administered to rabbits].
Apolipoprotein B plazmennyykh lipoproteidov, vstroennyi v liposomu: immunologicheskie svoistva i raspredelenie mezhdu organami pro vvedenii kroliku.

AB Apolipoprotein B (**apo B**) isolated from low density lipoproteins (**LDL**) was built in phospholipid-cholesterol liposomes, with the lipid/protein ratio being equal to 33:1. Such liposomes preserved their integrity, whereas the constituent **apo B** retained its antigenic properties. After intravenous injection to rabbits the pattern of **apo B liposome** distribution among organs was similar to that of **LDL**.
Apo B liposomes may be used for goal-oriented transport of some substances to organs and tissues whose cells have specific receptors for **apo B**-containing lipoproteins.

ACCESSION NUMBER: 84025000 MEDLINE
DOCUMENT NUMBER: PubMed ID: 6615606
TITLE: [Apolipoprotein B of plasma lipoproteins incorporated in liposomes: immunological properties and organ distribution when administered to rabbits].
Apolipoprotein B plazmennyykh lipoproteidov, vstroennyi v liposomu: immunologicheskie svoistva i raspredelenie mezhdu organami pro vvedenii kroliku.
AUTHOR: Klimov A N; Korovkin B F; Kuznetsov A S; Popov I N
SOURCE: Biulleten' eksperimental'noi biologii i meditsiny, (1983 Oct) 96 (10) 47-50.

Journal code: 0370627. ISSN: 0365-9615.

PUB. COUNTRY: USSR
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: Russian
FILE SEGMENT: Priority Journals
ENTRY MONTH: 198312
ENTRY DATE: Entered STN: 19900319
Last Updated on STN: 19900319
Entered Medline: 19831217

L2 ANSWER 5 OF 11 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
TI Anti-leishmanial drug delivery: Acetylated **LDL** as a site-specific delivery ligand.

AB The potential utility of acetylated **LDL** incorporated reverse-phase evaporation vesicles as J774.E1 macrophage specific delivery system was studied using Pentostam as anti-leishmanial drug and Leishmania mexicana mexicana as model macrophage disease. The investigations have shown that Apoprotein-B moiety of acetylated-**LDL** is incorporated into reverse-phase evaporation vesicles (acetylated-liposomes) allowing exploitation of the targeting properties of apoprotein-B ligand. Incorporation of **apo-B** into **liposome** carriers significantly enhances their uptake by Leishmania infected macrophages via the **LDL** and acetylated **LDL** receptors. The leishmanicidal action of Pentostam entrapped in acetylated liposomes was greater than native **LDL** containing liposomes and significantly higher than untargeted liposomes. Indeed targeted liposomes with acetylated **LDL** ligand have highly beneficial effect on the anti-leishmanial action of entrapped drugs and could contribute to a reduction in toxicity and increase in therapeutic index of currently prescribed anti-leishmanial drugs.

ACCESSION NUMBER: 2004:385437 BIOSIS
DOCUMENT NUMBER: PREV200400385943
TITLE: Anti-leishmanial drug delivery: Acetylated **LDL** as a site-specific delivery ligand.
AUTHOR(S): Shah, Akram [Reprint Author]; Hart, David
CORPORATE SOURCE: Dept ZoolParasitol Sect, Univ Peshawar, Peshawar, 25120, Pakistan
akramkokab@yahoo.com
SOURCE: Pakistan Journal of Zoology, (2004) Vol. 36, No. 1, pp. 45-52. print.
CODEN: PJZOAN. ISSN: 0030-9923.
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 29 Sep 2004
Last Updated on STN: 29 Sep 2004

L2 ANSWER 6 OF 11 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
TI Methylglyoxal and glyoxal enhance **LDL** oxidation through modification of **apo B** arginyl residues.

ACCESSION NUMBER: 2001:103429 BIOSIS
DOCUMENT NUMBER: PREV200100103429
TITLE: Methylglyoxal and glyoxal enhance **LDL** oxidation through modification of **apo B** arginyl residues.
AUTHOR(S): Mowri, Hiro-Omi [Reprint author]; Keaney, John F.
CORPORATE SOURCE: Boston Univ, Boston, MA, USA
SOURCE: Circulation, (October 31, 2000) Vol. 102, No. 18 Supplement, pp. II.82. print.
Meeting Info.: Abstracts from American Heart Association Scientific Sessions 2000. New Orleans, Louisiana, USA. November 12-15, 2000. American Heart Association.
CODEN: CIRCAZ. ISSN: 0009-7322.
DOCUMENT TYPE: Conference; (Meeting)

Conference; Abstract; (Meeting Abstract)

LANGUAGE: English

ENTRY DATE: Entered STN: 28 Feb 2001

Last Updated on STN: 15 Feb 2002

L2 ANSWER 7 OF 11 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

TI Oxidative interaction of unpaired hemoglobin chains with lipids and proteins: A key for modified serum lipoproteins in Thalassemia.

AB We searched for a biochemical explanation to the modification of lipoproteins like low-density lipoproteins (**LDL**) observed in patients with the severe hemolytic anemia beta-thalassemia. Because a large fraction of the **LDL** surface is composed of phospholipids, we first explored the possible involvement of phospholipids in the oxidative interaction of **LDL** with hemoglobin (Hb), using brain extract phospholipid liposomes as a model. The relative binding affinity and oxidative interaction of three hemoglobin variants (intact Hb A and isolated beta- and alpha-chains) with **LDL** and **liposome** were compared. Studies carried out at low pH/ionic strength and under physiological conditions revealed that association of hemoglobin variants with the phospholipid liposomes is driven by electrostatic forces but their binding is not a prerequisite for oxidative interaction. Unlike phospholipid liposomes, **LDL** underwent only a negligible association with the Hb variants under all pH/ionic strength conditions. Nevertheless, **LDL** induced oxidation of Hb variants, mostly alpha-chains. The dissimilar behavior of the liposomes and **LDL** indicated that **LDL** protein **apo B** rather than phospholipids is the actual **LDL** surface component which interacts with the hemoglobin variants. This agrees with the finding that **apo B** protein underwent oxidative crosslinking by the hemoglobin variants among which alpha-chains were most active. We concluded from these results that the ability of hemoglobin to undergo autooxidation is the key to its oxidative reactivity toward **LDL**. The results of the present study indicate that the modified **LDL** particles observed in beta-thalassemia may reflect lipoprotein oxidation by alpha-chains in circulation.

ACCESSION NUMBER: 1997:456085 BIOSIS

DOCUMENT NUMBER: PREV199799755288

TITLE: Oxidative interaction of unpaired hemoglobin chains with lipids and proteins: A key for modified serum lipoproteins in Thalassemia.

AUTHOR(S): Altamentova, Svetlana M.; Marva, Esther; Shaklai, Nurith [Reprint author]

CORPORATE SOURCE: Sackler Inst. Mol. Med., Sackler Fac. Med., Tel-Aviv Univ., Tel-Aviv 698887, Israel

SOURCE: Archives of Biochemistry and Biophysics, (1997) Vol. 345, No. 1, pp. 39-46.
CODEN: ABBIA4. ISSN: 0003-9861.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 27 Oct 1997

Last Updated on STN: 27 Oct 1997

L2 ANSWER 8 OF 11 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

TI LOW-DENSITY LIPOPROTEINS INTERACT WITH **LIPOSOME**-BINDING SITES ON THE CELL SURFACE.

AB Under physiological conditions significant amount of low-density lipoprotein **LDL** particles are taken up by cells independently of specific high-affinity **LDL** receptors (**apo-B** receptors). Previously it was established that some cells contain surface sites capable of binding liposomes. We proposed that **liposome**-binding sites could contribute to **LDL** interaction with the cell surface via phospholipid molecules of **LDL** particles. To check this hypothesis we studied the competitive interaction of human

LDL and DPPC liposomes with mouse embryo fibroblasts depleted of **apo-B** receptors by preliminary incubation with **LDL**. We have found that after removal of the **liposome** -binding sites from cell lamellae these areas of the cell surface lose their ability to bind **LDL**.

ACCESSION NUMBER: 1991:452391 BIOSIS
DOCUMENT NUMBER: PREV199192097171; BA92:97171
TITLE: LOW-DENSITY LIPOPROTEINS INTERACT WITH **LIPOSOME**
-BINDING SITES ON THE CELL SURFACE.
AUTHOR(S): GALKINA S I [Reprint author]; IVANOV V V; PREOBRAZHENSKY S
N; MARGOLIS L B; BERGELSON L D
CORPORATE SOURCE: BELOZERSKY LAB MOLECULAR BIOL BIOORGANIC CHEM, MOSCOW STATE
UNIV, MOSCOW 119899, MOSCOW, USSR
SOURCE: Febs Letters, (1991) Vol. 287, No. 1-2, pp. 19-22.
CODEN: FEBLAL. ISSN: 0014-5793.
DOCUMENT TYPE: Article
FILE SEGMENT: BA
LANGUAGE: ENGLISH
ENTRY DATE: Entered STN: 11 Oct 1991
Last Updated on STN: 11 Oct 1991

L2 ANSWER 9 OF 11 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
TI APO LIPO PROTEIN B OF PLASMA LIPO PROTEINS BUILT IN LIPOSOMES IMMUNOLOGIC
PROPERTIES AND DISTRIBUTION AMONG ORGANS AFTER ADMINISTRATION TO RABBITS.
AB Apolipoprotein B (**apo B**) isolated from low-density
lipoproteins (**LDL**) was built in phospholipid-cholesterol
liposomes, with a lipid/protein ratio of 33:1. Such liposomes preserved
their integrity; the constituent **apo B** retained its
antigenic properties. After i.v. injection to rabbits, the pattern of
apo B liposome distribution among organs was
similar to that of **LDL**. **Apo B** liposomes may
be used for goal-oriented transport of some substances to organs and
tissues whose cells have specific receptors for **apo B**
-containing lipoproteins.

ACCESSION NUMBER: 1984:291988 BIOSIS
DOCUMENT NUMBER: PREV198478028468; BA78:28468
TITLE: APO LIPO PROTEIN B OF PLASMA LIPO PROTEINS BUILT IN
LIPOSOMES IMMUNOLOGIC PROPERTIES AND DISTRIBUTION AMONG
ORGANS AFTER ADMINISTRATION TO RABBITS.
AUTHOR(S): KLIMOV A N [Reprint author]; KOROVKIN B F; KUZNETSOV A S;
POPOV I N
CORPORATE SOURCE: DEP BIOCHEM, INST EXP MED, ACAD MED SCI USSR, LENINGRAD,
USSR
SOURCE: Byulleten' Eksperimental'noi Biologii i Meditsiny, (1983)
Vol. 96, No. 10, pp. 47-50.
CODEN: BEBMAE. ISSN: 0365-9615.
DOCUMENT TYPE: Article
FILE SEGMENT: BA
LANGUAGE: RUSSIAN

L2 ANSWER 10 OF 11 JICST-EPlus COPYRIGHT 2005 JST on STN
TI Research and development of new medical imaging. 1.9. Research on the
localization change of HDL (Apo-A1) and **LDL (Apo-**
B) in vessel wall and arteriosclerosis lesion.
AB 1) In order to observe of peroxy lipid, formation process of the peculiar
autofluorescent material structural change of diabetes mellitus and
vascular change with fluorescent concentration-change was observed, and
the more detailed change of diabetic vascular change was recognized. 2)
From composition experiment by **liposome** model membrane and
culture smooth muscle cell, there were cell membrane failure and
structural change which showed photochemical reaction, and sensitivity
increase attributed to the hematoporphyrin which occurred here was
clarified.

ACCESSION NUMBER: 1040727302 JICST-EPlus
 TITLE: Research and development of new medical imaging. 1.9.
 Research on the localization change of HDL (Apo-A1) and
 LDL (Apo-B) in vessel wall and
 arteriosclerosis lesion.
 AUTHOR: MACHIDA MIKI
 CORPORATE SOURCE: Nippon Med. Sch.
 SOURCE: Nihon Ika Daigaku Haiteku Risachi Senta Kenkyu Hokoku
 Heisei 10nen 4gatsu- Heisei 15nen 3gatsu, (2003) pp. 128.
 Journal Code: N20041813
 PUB. COUNTRY: Japan
 DOCUMENT TYPE: Journal; Article
 LANGUAGE: Japanese
 STATUS: New

L2 ANSWER 11 OF 11 WPIDS COPYRIGHT 2005 THE THOMSON CORP on STN
 TI Treatment of angina or angina equivalent, e.g. dyspnea, arrhythmia by
 administering large liposomes comprised of phospholipids substantially
 free of sterols.
 AN 2001-080232 [09] WPIDS
 AB WO 200069412 A UPAB: 20050211
 NOVELTY - A method for treating angina or anginal equivalent comprises
 administering a multiplicity of large liposomes comprised of phospholipids
 substantially free of sterols is new.
 DETAILED DESCRIPTION - A method for treating angina comprises
 administering a multiplicity of large liposomes comprised of phospholipids
 substantially free of sterols is new.
 INDEPENDENT CLAIMS are also included for
 (1) a pharmaceutical kit for treating angina or anginal equivalent
 comprising a first container having liposomes; and a second container
 having anti-anginal drugs other than said liposomes;
 (2) a method of perioperative and/ or pre-operative conditioning of a
 subject comprising liposomes; and
 (3) a method for treating claudication.
 ACTIVITY - Cardiant; antiarrhythmic; antianginal; hypotensive;
 analgesic; antianemic.
 MECHANISM OF ACTION - Angiotensin-converting enzyme(ACE) inhibitor.
 USE - To monitor a level of plasma atherogenic lipoprotein, a cardiac
 function preferably EKG abnormality, an S-T segment change, arrhythmia, an
 assessment of segmental wall motion, blood viscosity, exercise tolerance,
 ambulatory EKG monitoring and to treat angina, preferably stable angina,
 unstable angina, variant angina, angina pectoris, an anginal equivalent
 selected from an ischemic wall motion abnormality, dyspnea, impaired
 exercise tolerance, an arrhythmia, a reduced cardiac function, shortness
 of breath, fatigue, abdominal distress and referred pain, hypertension,
 hyperthyroidism, pulmonary disease, heart failure, hypermetabolic state,
 anemia and claudication.(all claimed)
 ADVANTAGE - The combination with drugs enhances the intracellular
 movement of cholesterol to the cell membrane.
 Dwg.0/28

ACCESSION NUMBER: 2001-080232 [09] WPIDS
 DOC. NO. CPI: C2001-022961
 TITLE: Treatment of angina or angina equivalent, e.g. dyspnea,
 arrhythmia by administering large liposomes comprised of
 phospholipids substantially free of sterols.
 DERWENT CLASS: B05
 INVENTOR(S): GOLDBERG, D; WILLIAMS, K J; GOLDBERG, D I
 PATENT ASSIGNEE(S): (TALA-N) TALARIA THERAPEUTICS INC; (ESPE-N) ESPERION LUV
 DEV INC; (ESPE-N) ESPERION MERGERCO INC
 COUNTRY COUNT: 91
 PATENT INFORMATION:

PATENT NO	KIND DATE	WEEK	LA	PG
-----------	-----------	------	----	----

```

-----
WO 2000069412  A1 20001123 (200109)* EN 142
RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW NL
   OA PT SD SE SL SZ TZ UG ZW
W:  AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES
   FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS
   LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL
   TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW
AU 2000050053  A 20001205 (200113)
EP 1183011     A1 20020306 (200224) EN
R:  AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT
   RO SE SI
JP 2003508349  W 20030304 (200319)      184
AU 773385      B2 20040527 (200465)
AU 2004203419  A1 20040819 (200474)#
AU 2004203419  A2 20040819 (200510)

```

APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
WO 2000069412	A1	WO 2000-US12962	20000512
AU 2000050053	A	AU 2000-50053	20000512
EP 1183011	A1	EP 2000-932314	20000512
		WO 2000-US12962	20000512
JP 2003508349	W	JP 2000-617871	20000512
		WO 2000-US12962	20000512
AU 773385	B2	AU 2000-50053	20000512
AU 2004203419	A1	AU 2004-203419	20040727
AU 2004203419	A2	AU 2004-203419	20040727

FILING DETAILS:

PATENT NO	KIND	PATENT NO
AU 2000050053	A Based on	WO 2000069412
EP 1183011	A1 Based on	WO 2000069412
JP 2003508349	W Based on	WO 2000069412
AU 773385	B2 Previous Publ. Based on	AU 2000050053 WO 2000069412
AU 2004203419	A1 Div ex	AU 773385
AU 2004203419	A2 Div ex	AU 773385

PRIORITY APPLN. INFO: US 1999-134140P 19990514; AU
2004-203419 20040727

=> e halbert, g,au

```

E1      1      HALBERSTSMA/BI
E2     34      HALBERT/BI
E3      0 --> HALBERT, G,AU/BI
E4      3      HALBERTI/BI
E5      4      HALBERTSMA/BI
E6      1      HALBERTSTADT/BI
E7      1      HALBERTSTAEDTER/BI
E8      3      HALBERZEUGNISSE/BI
E9      5      HALBES/BI
E10     1      HALBESBERG/BI
E11     1      HALBESTERN/BI
E12     3      HALBETASOL/BI

```

=> e halbert, g/au

```

E1      3      HALBERT V A/AU

```

E2	2	HALBERT W M/AU
E3	0 -->	HALBERT, G/AU
E4	2	HALBERTAL E/AU
E5	20	HALBERTHAL E/AU
E6	9	HALBERTHAL M/AU
E7	1	HALBERTHAL M S/AU
E8	7	HALBERTHAL MICHAEL/AU
E9	1	HALBERTHAL MICHAEL S/AU
E10	1	HALBERTHAL MIKI/AU
E11	1	HALBERTHAL R J/AU
E12	2	HALBERTLAS R/AU